

Fighting H1N1 swine flu is a three-step approach: 1) get a vaccination when it becomes available, 2) practice everyday prevention (including frequent hand washing, covering coughs and sneezes and staying home when sick), and 3) use antiviral drugs correctly if your health professional recommends them.

How do I know if I have the flu?

You may have the flu if you have some or all of these symptoms:

- fever *
- cough
- sore throat
- runny or stuffy nose
- body aches
- headache
- chills
- fatigue
- sometimes diarrhea and vomiting

*It's important to note that not everyone with flu will have a fever.

What should I do if I get sick?

If you get sick with flu-like symptoms this flu season, you should stay home and avoid contact with other people except to get medical care. Most people with [2009 H1N1](#) have had mild illness and have not needed medical care or antiviral drugs and the same is true of seasonal flu.

However, some people are more likely to get flu complications and they should talk to a health care provider about whether they need to be examined if they get flu symptoms this season. They are:

- Children younger than 5, but especially children younger than 2 years old
- People 65 and older
- Pregnant women
- People who have: Chronic Illnesses.

Also, it's possible for healthy people to develop severe illness from the flu so anyone concerned about their illness should consult a health care provider.

There are emergency warning signs. Anyone who has them should get medical care right away.

What are the emergency warning signs?

In children

- Fast breathing or trouble breathing
- Bluish skin color
- Not drinking enough fluids
- Not waking up or not interacting
- Being so irritable that the child does not want to be held
- Flu-like symptoms improve but then return with fever and worse cough
- Fever with a rash

In adults

- Difficulty breathing or shortness of breath
- Pain or pressure in the chest or abdomen
- Sudden dizziness
- Confusion
- Severe or persistent vomiting

How long should I stay home if I'm sick?

[CDC recommends that you stay home for at least 24 hours](#) after your fever is gone except to get medical care or for other things you have to do and no one else can do for you. (Your fever should be gone without the use of a fever-reducing medicine, such as Tylenol®.) You should stay home from work, school, travel, shopping, social events, and public gatherings.

What should I do while I'm sick?

Stay away from others as much as possible to keep from making them sick. If you must leave home, for example to get medical care, wear a facemask if you have one, or cover coughs and sneezes with a tissue. And wash your hands often to keep from spreading flu to others. CDC has information on "[Taking Care of a Sick Person in Your Home](#)" on its website at http://www.cdc.gov/h1n1flu/guidance_homecare.htm